

- target genes of the bZIP transcription factor OsTGAP1, whose overexpression causes elicitor-induced hyperaccumulation of diterpenoid phytoalexins in rice cell. PLOS ONE, 9, e105823.
- Miyamoto, K. et al. 2016. Evolutionary trajectory of phytoalexin biosynthetic gene clusters in rice. Plant J., 87, 293-304.
- Okada, A. et al. 2009. OsTGAP1, a bZIP transcription factor, coordinately regulates the inductive production of diterpenoid phytoalexins in rice. J. Biol. Chem., 284, 26510-26518.
- Okada, K. 2011. The biosynthesis of isoprenoids and the mechanisms regulating it in plants. Biosci. Biotechnol. Biochem., 75, 1219-1225.
- Otomo, K. et al., 2004. Biological functions of ent- and syn-copalyl diphosphate synthases in rice: key enzymes for the branch point of gibberellin and phytoalexin biosynthesis. Plant J., 39, 886-893.
- Peters, R. J. 2006. Uncovering the complex metabolic network underlying diterpenoid phytoalexin biosynthesis in rice and other cereal crop plants. Phytochemistry, 67, 2307-2317.
- Sakamoto, T. et al. 2004. An overview of gibberellin metabolism enzyme genes and their related mutants in rice. Plant Physiol., 134, 1642-1653.
- Sekido, H. et al. 1986. Oryzalexin D (3,7-dihydroxy-(+)-sandaracopimaradiene), a new phytoalexin isolated from blast infected rice leaves. J. Pestic. Sci., 11, 369-372.
- Shimura, K. et al. 2007. Identification of a biosynthetic gene cluster in rice for momilactones. J. Biol. Chem., 282, 34013-34018.
- Tamogami, S. et al. 1993. Oryzalexin S structure: a new stearane-type rice plant phytoalexin and its biogenesis. Tetrahedron, 49, 2025-2032.
- Toyomasu, T. 2008. Recent advances regarding diterpene cyclase genes in higher plants and fungi. Biosci. Biotechnol. Biochem. 72, 1168-1175.
- Toyomasu, T. et al. 2008. Diterpene phytoalexins are biosynthesized in and exuded from roots of rice seedlings. Biosci. Biotechnol. Biochem., 72, 562-567.
- Toyomasu, T. and Sassa, T. 2010. Comprehensive Natural Products II Chemistry and Biology, Vol. 1. Oxford: Elsevier, pp. 643-672.
- Toyomasu, T. et al. 2014. Reverse-genetic approach to verify physiological roles of rice phytoalexins: characterization of a knockdown mutant of *OsCPS4* phytoalexin biosynthetic gene in rice. Physiol. Plant. 150, 55-61.
- Toyomasu, T. et al. 2015. Transcripts of two *ent*-copalyl diphosphate synthase genes differentially localize in rice plants according to their distinct biological roles. J. Exp. Bot. 66, 369-376.
- Wu, Y. et al. 2012. Functional characterization of wheat copalyl diphosphate synthases sheds light on the early evolution of labdane-related diterpenoid metabolism in the cereals. Phytochemistry, 84, 40-46.
- Xu, M. et al. 2012. Genetic evidence for natural product-mediated plant-plant allelopathy in rice (*Oryza sativa*). New Phytol., 193, 570-575.
- Yajima, A. and Mori, K. 2000. Diterpenoid total synthesis. XXXII. Synthesis and absolute configuration of (-)-phytocassane D, a diterpene phytoalexin isolated from the rice plant, *Oryza sativa*. Eur. J. Org. Chem., 2000, 4079-4091.
- Yamaguchi S. 2008. Gibberellin metabolism and its regulation. Ann. Rev. Plant Biol., 59, 225-251.
- Yamamura, C. et al. 2015. Diterpenoid phytoalexin factor, a bHLH transcription factor, plays a central role in the biosynthesis of diterpenoid phytoalexins in rice. Plant J., 84, 1100-1113.
- Yokotani, N. et al. 2013. OsWRKY76 is a rice transcriptional repressor playing opposite roles in blast disease resistance and cold stress tolerance. J. Exp. Bot., 64, 5085-5097.
- Zhou, K. et al. 2012. Functional characterization of wheat *ent*-kaurene(-like) synthases indicates continuing evolution of labdane-related diterpenoid metabolism in the cereals. Phytochemistry, 84, 47-55.